

Vol-12 Issue-01 April 2023

Weaned pigs' health, oxidative stress state, and fatty acid profile as a result of sodium humate and zinc oxide used for post-weaning diarrhea prevention

S.K. JATA, M. NEDUNCHEZHIYAN

ABSTRACT: In this study, weaned piglets were tested for oxidative stress, serum fatty acid (FA) profile, and clinical, biochemical, and haematological markers of health status in relation to sodium humate (HNa), which was used as a partial replacement for pharmacological doses of zinc oxide (ZnO) in feed or drinking water. The 32 piglets that had just been weaned were divided into four groups: The control group received a baseline meal containing 110 mg ZnO/kg. The half-dosage group received a pharmacological dose of 2.5 g ZnO/kg. The full-dose group received 1.7 g ZnO and 20.0 g HNa/kg. The water group received 1.7 g ZnO/kg along with 0.2% HNa. Unlike the Control group, which included three piglets who were sick with diarrhea, the piglets treated with ZnO showed no adverse effects and performed well clinically. As compared to the Control group, those treated with ZnO 2.5 or ZnO 1.7 + HNa (w) had higher levels of triacylglycerols, suggesting an improvement in energy metabolism. Treatments had no effect on total cholesterol, HDL, or LDL. Serum urea levels were lower in piglets given ZnO and HNa (ZnO 1.7 + HNa (w)) or showed a tendency toward lower levels (ZnO 1.7 + HNa (f)) for both treatments. The blood of the treated pigs showed considerably greater haematocrit and hemoglobin levels in the ZnO 2.5 and ZnO 1.7 + HNa (f) treatments, but no significant difference in the ZnO 1.7 + HNa (w) treatments. An rise in blood 8-iso-PGF2a, the most dependable indicator of oxidative stress, was seen after long-term ZnO 2.5 therapy. As shown by substantial (ZnO 1.7 + HNa (w)) or non-significant (ZnO 1.7 + HNa (f)) decreases in serum 8-iso-PGF2 α , the oxidative state of pigs was favorably impacted by partial replacement of ZnO with HNa. Serum levels of saturated (SFA), monounsaturated (MUFA), and n-3 polyunsaturated FA (PUFA) were all markedly reduced after treatment with ZnO 2.5, whereas levels of PUFA n-6 and n-6/n-3 ratios were markedly raised, in comparison to the Control group. The production of n-6 dihomo-y-linolenic acid and n-3 eicosapentaenoic and docosapenoic longchain polyunsaturated fatty acids (LC-PUFA) was adversely impacted by ZnO 2.5 treatment. The percentage of SFA and PUFA n-3 was considerably raised in both the ZnO 1.7 + HNa (f)/(w) and ZnO 2.5 treatments, whereas the proportion of PUFA n-6 and the n-6/n-3 ratios were lowered. The impact was more pronounced in the ZnO 1.7 + HNa (w) treatment. Both treatments were quite comparable to Control in terms of individual PUFA n-3 and n-6 and the n-6/n-3 ratio. Compared to a high pharmacological dosage of ZnO, the effects of HNa as a partial ZnO substitute on weaned piglets' performance and health are comparable, and it may mitigate the oxidative stress caused by long-term ZnO over-supplementation. Plus, this therapy may reverse the negative effects of high ZnO dosages on the n-6/n-3 ratio and blood concentrations of some physiologically important LC-PUFA. In general, it is safe to say that adding HNa to both feed and water has comparable effects, although the effects on drinking water are noticeably stronger.

Keywords: subject areas include pigs, lipid metabolism, eicosanoids, performance, serum biochemistry, haematology, and gas chromatography/mass spectrum



INTRODUCTION:

AA= arachidonic acid, ALA = α -linolenic acid, ALP = alkaline phosphatase, ALT = alanine transaminase, AST = aspartate aminotransferase, BWG = body weight gain, DGLA = dihomo- γ -linolenic acid, DHA = docosahexae- noic acid, DPA = docosapentaenoic acid, EPA = eicosapentaenoic acid, ETEC = enterotoxigenic *Escherichia coli*, FA = fatty acids, FAME = fatty acid methyl esters, FCR = feed conversion ratio, FI = feed intake, GC/MS = gas Supported by the Ministry of Agriculture of the Czech Republic (Grants No. QJ1210112 and No. RO0516) and by the Ministry of Education, Youth and Sports of the Czech Republic under the NPU I Program (Grant No. LO1218).

doi: 10.17221/70/2016-VETMED

chromatography/mass spectrometry, **HA** = humic acid, **HCT** = haematocrit, **HDL** = high-density lipoprotein, **HGB** = haemoglobin, **HNa** = sodium humate, **HPLC** = high performance liquid chromatography, **HS** = humic substances, **LA** = linoleic acid, **LC-MS/MS** = liquid chromatography/tandem mass spectrometry, **LC-PUFA** = long-chain polyunsaturated fatty acid, **LDL** = low-density lipoprotein, **MUFA** = monounsaturated fatty acids, **OA** = oleic acid, **PA** = palmitic acid, **PUFA** = polyunsaturated fatty acids, **ROS** = reactive oxygen species, **SFA** = saturated fatty acids, **TGC** = triacylglycerols, **USFA** = unsaturated fatty acids

The beneficial effects of pharmacological doses of zinc oxide (ZnO) in piglets in the postweaning period are well documented (Kim et al. 2012; Heo et al. 2013). In the last decades, the widespread use of high levels of ZnO has been criticised because of the subsequent high excretion of Zn into the environment. EFSA limits the maximum contentof ZnO in complete feeding stuffs to 250 mg/kgfor all animal species (EC Regulation 1831/2003), although the use of higher pharmacological dosesis still permitted under veterinary prescription as the need arises. There are ongoing efforts aimedat finding effective alternatives, that could protect piglets from post-weaning diarrhoea and growth depression after weaning. In our previous studies (Trckova et al. 2015; Kaevska et al. 2016) we veri-fied the possibility of the substitution of a pharmacological dose of ZnO (2.5 g/kg diet) by humic substances (HS). HS are natural bioactive agents primarily decomposed from organic matter by bacteria in the soil. They have been reported to improve performance and

ISSN: 2320-3730

Vol-12 Issue-01 April 2023

nutrient utilisation and show anti-bacterial, antiviral and anti-inflamma- tory effects (Islam et al. 2005; Ji et al. 2006; Wang et al. 2008). EMEA (1999) has recommended humic acid (HA) and its sodium salt for the treatment of diarrhoea, dyspepsia and acute intoxications in animals. However, our previous results showed that in cases of severe post-weaning diarrhoeal infec- tions caused by enterotoxigenic Escherichia coli strains (ETEC), supplementation of HS to piglets can be adequately effective only in combination with a specific dose of ZnO (1.7 g/kg; Trckova et al. 2015). Moreover, there is still a lack of data about other potential effects of this partial substitution of ZnO by HS on the young and vulnerable organism of weaning piglets. Weaning is the most difficult period in a piglet's life and is associated with social stress and dietary changes. These can lead to digestive disorders, which cause diarrhoea, and increased disease susceptibilityresulting in economic losses in the post-weaning period (Metzler et al. 2005; Heo et al. 2013). Weaning stress, especially in early-weaned pigs, can also re-sult in the suppression of antioxidant enzyme activ-ity as well as in promotion of free radical production leading to the oxidative damage of lipids, proteins, and DNA (Zhu et al. 2012; Yin et al. 2014). This oxi-dative imbalance can result in immunosuppression with increased susceptibility to various diseases and in negative alterations in the growth performance of animals. Oxidative stress may also have a negative effect on meat quality and the shelf life of meat(Aksu et al. 2005; Bai et al. 2013). In some reports, it was remarked that HS may have the potential to negate the effect of oxidative stress within the body (Ipek et al. 2008; Weber et al. 2014). On the other hand, HS, specifically HA, administered over a long period at high levels has been reported to induce damage in cells and tissues accompanied by an increase in oxidative stress andlipid peroxidation (Cheng et al. 1999; Varanka et al. 1999; Gau et al. 2001; Ho et al. 2003; Qi et al. 2008). HS are natural materials with variable composition, which may explain the varying efficacy described in the literature.



HS have been reported to influence saturated (SFA) and unsaturated fatty acid (USFA) concentrations and their ratios in the meat of finishing pigs (Wanget al. 2008). Nutrition, biosynthesis, lipogenesis and remodelling are the main sources of fatty acids (FA) and lipid composition in piglets. However, it can be supposed that the metabolism and biosynthesis of FA and lipids in young growing piglets could also be regulated in a different manner. The aim of this study was to investigate the effects of sodium humate (HNa), applied to feed or drinking water as a partial substitution for phar- macological doses of ZnO, on clinical, biochemical and haematological indicators of health status, a

biomarker of oxidative stress (serum 8-*iso*-PGF_{2 α}) and the serum FA profile in weaned piglets.

ATERIAL AND METHODS

Animal management and dietary treatment. Testing of HNa and ZnO was performed on 32 weaned piglets $[LW \times (P \times Du)]$ originating from aspecific pathogen-free herd. Piglets were transported to the experimental animal facility of the Veterinary Research Institute, Brno, Czech Republic on the day of weaning (28th day of age). They were identified by individual ear tags and housed in indoor pens. Animal handling followed EU directive 86/609/EEC concerning animal care. The animal care protocol for this experiment followed the Czech guidelines for animal experimentation and was approved by the Branch Commission for Animal Welfare of the Ministry of Agriculture of the Czech Republic (Permission No. MZe 50-2011).

Weaned piglets with an average initial body weight of 8.79 ± 1.31 kg were allocated into Ingredient (g/kg) Basal diet Wheat Barley 298.0 Soybean meal (47% CP) 186.0 Dry whey and soy protein concentrate 50.0 Soybean oil 23.0 Limestone, ground12.0 Dicalcium phosphate 11.0 Salt 3.0 Sodium carbonate 1.0 **l-Lysine HCl** 0.2 1-Threonine 0.3 dl-Methionine 0.3 Vitamin and mineral premix¹ 15.0 Calculated chemical composition

ISSN: 2320-3730

Vol-12 Issue-01 April 2023

four groups. Each group was kept in one pen with eight piglets (with an equal sex ratio of four males and four females) per pen. Dietary treatments were as follows: Control = basal diet containing 110 mg ZnO/kg; ZnO 2.5 = basal diet supplemented with a pharmacological dose of ZnO (2.5 g ZnO/kg); ZnO 1.7 + HNa (f) = basal diet supplemented with

1.7 g ZnO and 20.0 g HNa/kg; ZnO 1.7 + HNa

(w) = basal diet supplemented with 1.7 gZnO/kg and drinking water supplemented with 0.2% HNa. The concentration of HNa in the drinking water was chosen so that the average daily intake of HNaper piglet by water was of approximately the same dose as the average daily intake of HNa by diet. The basal diet (Table 1) was formulated accord-ing to animal requirements (NRC 1998). HNa was made from oxyhumolite (Humatex, a.s., Bilina, Czech Republic) by neutralisation and extraction of the present humic acids with sodium hydroxide in aqueous medium. The chemical analysis revealed 93.9% dry matter, 34.3% ash, 52.4% HS, 15.2% HA and 37.2% fulvic acid in dry matter. Piglets were fed twice a day *ad libitum* and water was provided by automatic waterers. The dietary treatment was maintained for 21 days.

Clinical, biochemical and haematological indicators of health status. Piglets were observed daily for morbidity and mortality. Faecal consistency was recorded daily. To assess the general health status of piglets after long-term treatments, 10 ml samples of blood were drawn from the *vena cava cranialis* after 12-h fasting at the end of the trial

Table 1. Composition of the diet for weaned piglets (as-fed basis)

400.2



ME (MJ/kg)	12.27
Crude protein (g/kg)	189.00
Fat (g/kg)	4.10
Lysine (g/kg)	11.95
Methionine (g/kg)	4.00
Ca (g/kg)	7.00
Na (g/kg)	1.90

Vol-12 Issue-01 April 2023



P (g/kg) 4.55

Vol-12 Issue-01 April 2023

¹Provided per kg diet: 12 000 IU of vitamin A, 2000 IU of vitamin D3, 100 IU of vitamin E, 152 mg of Cu (as CuSO₄), 22 mg of Zn (as ZnO), 88 mg of Zn (as ZnSO₄), 32 mg of Mn (as MnO), 110 mg of Fe (as FeSO₄), 1.0 mg of I (as Ca(IO₃)₂), 0.20 mg of Co (as Co₂O₃ × 7 H₂O), and 0.3 mg of Se (as

$Na_2SeO_3 \times 5 H_2O)$

for monitoring of the haematological and serum biochemical profile. From blood samples collected into heparinised tubes, haematocrit (HCT), hae- moglobin (HGB), erythrocyte and leukocyte counts were determined using the Coulter Counter M4 ap-paratus (Coulter Cientifica S.A., Mostoles, Spain), whereas the differential leukocyte count in blood smears was determined using a Nikon Eclipse E600 fluorescence microscope (Nikon, Tokyo, Japan).

Serum was separated from blood samples by cen- trifugation at 4000 rpm for 20 min and stored at -20 °C until further analyses. In serum samples, glucose, total protein, albumin, triacylglycerol (TGC), cholesterol, high-density lipoprotein (HDL) and low-density lipoprotein (LDL), aspartate ami- notransferase (AST), alanine transaminase (ALT), alkaline phosphatase (ALP), calcium, phosphorus, doi: 10.17221/70/2016-VETMED

magnesium, iron and zinc were measured using a BS200 automated chemistry analyser (Mindray, Shenzhen, China).

Piglets were weighed at the beginning (Day 1) and at the end (Day 21) of the trial. Individual body weight gains (BWG) were calculated. The average daily feed intake (FI) of the groups was recorded. The feed conversion ratio (FCR) was calculated from the FI and BWG.

Biomarker of oxidative stress. Solid phase ex- traction was used for extraction of 8-*iso*-PGF_{2 α} from blood serum. Select HLB SPE (1 ml, 30 mg)

cartridges (Supelco, Prague, Czech Republic) were washed with 1 ml of ethyl acetate, 1 ml of metha- nol and 1 ml of water. Then, 0.5 ml of serum were loaded onto the SPE column. The columns were then washed with 2 ml of water. The cartridges wereair dried for 3 min with vacuum and analytes were eluted with 1.5 ml of methanol. Samples were dried under a stream of nitrogen, dissolved in 60 μ l of methanol and aliquots of 5 μ l were injected into thehigh-performance liquid chromatography (HPLC) column. Analysis of samples was performed using liquid chromatography/tandem mass spectrometry (LC-MS/MS). An Agilent 1200 chromatographic sys- tem (Agilent Technologies, Germany) was used. Separation of prostaglandins was carried out using a HPLC Ascentis Express C18 column (2.1 × 150 mm, 2.7 μ m particle size, Supelco, Bellefonte, USA) maintained at 45 °C. Mobile phases consisted of 0.1% formic acid in water and acetonitrile. A 25-min linear gradient from 30 to 100% acetonitrile was used. The flow rate was 0.3 ml/min, and the in- jection volume was 5 μ l. The HPLC system was coupled to an Agilent 6410 Triple Quad LC/MS triple quadrupole mass spectrometer (Agilent Technologies, USA). Electrospray ionisation was performed in negative ion mode with nitrogen as

the nebulising and collision gas. 8-*iso*-PGF_{2 α} was analysed using multiple reaction monitoring and mass transition; precursor and product ions were 353.2/193.1 *m/z*.

FA profile in diets and blood serum. The blood serum was protected from oxidation by the addi- tion of butylated hydroxytoluene at a final con- centration of 40 mg/ml and was stored at -80 °C. Samples were analysed within one month of collec- tion. After protein precipitation, 0.5 ml of serum and about 1.0 g of diets and HNa were extracted according to a modified protocol of Bligh and Dyer

(1959). Appropriate volumes of extracts were evap- orated to dryness under a stream of nitrogen and a slightly modified method of Kang and Wang (2005) was used for the transesterification of FA from extracted lipids. Briefly, 20 μ l of the internal stand- ard solution (C23:0 FA, 80 μ g/ml), 20 μ l butylated hydroxytoluene (200 μ g/ml), 0.5 ml of hexane and

0.5 ml of 10% BF₃ in methanol were added to the drysample in a 4 ml silanised glass vial. Samples were vortexed, vials were capped under nitrogen andtransferred to a heating block at 100 °C and heated for 60 min. Subsequently, the samples were cooled, uncapped, and neutralised by the addition of 1 ml of a 5% solution of K_2CO_3 . The FA methyl esters (FAME) were then extracted twice with 1 ml of hex-



Vol-12 Issue-01 April 2023

ane. Combined extracts were evaporated to drynessunder a stream of nitrogen and dissolved in 200 μ l of 2,2,4-trimethylpentane. From the final samples, 1 μ l was injected into the capillary column of a gas chromatography/mass spectrometry (GC/MS) system.

GC/MS analysis was based on determination of retention times and relative abundances of selected mass ions of FAME. GC separation was done in anOmegawax fused silica capillary column (30 m \times 0.25 mm *i.d.*, 0.25 μ m, Supelco, Bellefonte, USA).

Helium at a column head pressure of 70 kPa was used as the carrier gas. An ion trap mass spectrometer Saturn 2100T (Varian, Walnut Creek, USA) was used for the detection and identification of analytes. The mass spectrometer was operated in electron ionisation and scan mode at an electron energy of 70 eV. Relative standard deviation for three parallel assays ranged from 1.8% (22:0) to 16.9% (20:4n3).

Individual FA were identified by n-designation (car-bon numbering starts from the methyl end of the molecule). A total of 50 FA were analysed. Only FA whose concentration was higher than 0.5 nmol/ml were summarised.

Statistical analyses. All data were subjected to statistical analysis using the GraphPad Prism 5.04 (GraphPad, Inc., San Diego, USA) and Statistica

12.6 (Dell, Inc., Tulsa, USA) software. The normal- ity of the data was tested with the D'Agostino and Pearson omnibus normality test and the homoge- neity of variances using the Brown-Forsythe test. When data passed the normality and homoscedas- ticity test, the statistical significance of the differences among the group means was determined by the analyses of variance (ANOVA) in conjunc-

tion with the Tukey's multiple comparison test. Otherwise, the Kruskal-Wallis test in conjunction with the Dunn's multiple comparison test were used. Mortality and diarrhoea incidence data were analysed using the Chi-squared test. Differences between means with P < 0.05 were accepted as be- ing statistically significant.

RESULTS AND DISCUSSION

Clinical, biochemical and haematologicalindicators of health status

All ZnO treatments, at pharmacological dosesor at amounts supplemented with HNa in feed or drinking water, resulted in good health status of weaned piglets. None of the piglets from the treated groups expressed any clinical signs of illness over the course of the experimental period. In the untreated control group, clinical signs of diarrhoea (watery faeces) were recorded in three out of the eight piglets. Diarrhoea occurred on the secondweek after weaning (Day 8) and lasted for four to five days in individual piglets.

Partial substitution of ZnO by HNa resulted in good performance, which was comparable to pharmacological dosing with ZnO (Table 2). The BWG and FI of treated groups did not differ significantly; significantly higher (P < 0.01) BWG values were detected only in comparison with the untreated Control. Piglets from treated groups were also more efficient in converting feed to gain than Control. Differences in performance after HNa supplemen-

Table 2. Performance and diarrhoea occurrence in pigletsdoi: 10.17221/70/2016-VETMED

tation in feed and water were not significant, al- though there was a trend towards better BWG and FCR values in ZnO 1.7 + HNa (w).

The results are in accordance with previous find- ings that confirmed the positive effect of ZnO and HNa treatments on performance in weaned piglets (Trckova et al. 2015; Kaevska et al. 2016). ZnO is well known to enhance growth performance in piglets through the stabilisation of the intestinal microbiota and improvements in nutrient diges- tion and utilisation. HS may benefit animal performance even though the actual mechanism is not fully understood (Kim et al. 2004; Ji et al. 2006; Bai et al. 2013).



Vol-12 Issue-01 April 2023

The high, pharmacological dose of ZnO as well as lower dose supplemented with HNa altered the serum chemistry of piglets, particularly the activity of liver enzymes (Table 3). Increases in liver enzyme activities after ZnO and HS dietary administration were reported previously (Poulsen 1995; Trckovaet al. 2015). However, the increased levels of liver enzymes in treated piglets were found to be within the physiological ranges (Jackson and Cockcroft 2002) and they did not reflect any trend that would suggest any adverse effect of treatments.

Apart from the activity of liver enzymes, TGC, urea, Ca and Zn were also significantly affected by treatments. Significantly increased TGC in ZnO

2.5 and ZnO 1.7 + HNa (w) in comparison with Control suggested increased energy metabolism in piglets after treatments. This is in accordance with Zhang et al. (2014), who reported a dose- dependent increase in TGC with supplementa- tion of dietary Zn. On the other hand, HS were

]	Treatments	-Main Pos	st-hoc tests								
Parameter HNa (f) HNa (w)	Control	ZnO 2.5	ZnO 1.7 +	ZnO 1.7 +	test	1:2	1:3	1:4	2:3	2:4	3:4
Initial body weight (kg)	8.78 ± 1.43	8.81 ± 1.47	8.80 ± 1.28	8.74 ± 1.32	ns						
Body weight gain (g/day)	191.5 ± 0.03	286.8 ± 0.03	249.5 ± 0.04	271.9 ± 0.04	**	赤赤	非非	非非	ns	ns	ns
Feed intake (g/pig/day)	453.5 ± 287.5	500.9 ± 327.5	495.8 ± 324.6	6 492.4 ± 304.8	ns						
Feed conversion ratio	2.37	1.75	1.99	1.81							
Mortality (piglets)	0	0	0	0	ns						
Diarrhoea incidence (piglet	s) 3	0	0	0	ns						
Duration of diarrhoea (days	s) 1.75	0	0	0	ns						

Control = basal diet containing 110 mg ZnO/kg, ZnO 2.5 = basal diet supplemented with 2.5 g ZnO/kg, ZnO 1.7 + HNa(f) = basal diet supplemented with 1.7 g ZnO and 20.0 g HNa/kg, ZnO 1.7 + HNa (w) = basal diet supplemented with 1.7 g ZnO/kg and drinking water supplemented with 0.2% HNa

P ₹0.01

i: 10.17221/70/2016-VETMED

Table 3. Serum biochemical and haematological parameters of piglets on Day 21 after weaning

Control ZnO 2.5 ^{ZnO 1.7} ZnO 1.7 +	+		HNa (f)								
HNa (w)test											
1:2 1:3 1:4 2:3 2:4 3:4											
Albumin (g/l)	26.45 ± 2.77	29.42 ± 1.66	28.57 ± 1.28	27.45 ± 2.40	ns						
Total protein (g/l)	44.59 ± 3.24	45.11 ± 4.87	44.97 ± 4.09	44.89 ± 5.63	ns						
Triacylglycerols (mmol/l)	0.28 ± 0.07	0.45 ± 0.06	0.30 ± 0.08	0.42 ± 0.11	₩₩	*	ns	*	*	ns	ns
Cholesterol (mmol/l)	2.05 ± 0.16	2.26 ± 0.19	2.14 ± 0.30	2.26±0.39	ns						
HDL (mmol/l)	1.33 ± 0.19	1.24 ± 0.10	1.16 ± 0.19	1.20 ± 0.21	ns						
LDL (mmol/l)	0.70 ± 0.11	0.86 ± 0.11	0.76±0.15	0.81 ± 0.17	ns						
Creatinine (µmol/l)	95.68 ± 15.34	92.98 ± 6.40	87.88 ± 10.68	92.09 ± 6.55	ns						
Urea (mmol/l)	4.25 ± 1.39	3.60 ± 0.89	2.86 ± 1.21	2.50 ± 0.93	*	ns	ns	*	ns	ns	ns
aLT (µkat/l)	0.78 ± 0.07	1.02 ± 0.09	0.98 ± 0.12	0.90 ± 0.15	察察	**	**	ns	ns	ns	ns
aST (µkat/l)	0.57 ± 0.09	0.62 ± 0.12	0.84 ± 0.11	0.72 ± 0.08	察察	ns	非非	*	非非	ns	ns
aLP (µkat/l)	3.59 ± 1.02	8.13 ± 1.56	5.43 ± 0.91	4.56 ± 1.07	察察	非非	÷	ns	非非	赤赤	ns
Ca (mmol/l)	2.73 ± 0.17	3.08 ± 0.20	2.84 ± 0.11	2.79 ± 0.19	₩₩	**	ns	ns	ns	*	ns
P (mmol/l)	2.78 ± 0.24	2.72 ± 0.21	2.86 ± 0.37	2.88±0.28	ns						
Mg (mmol/l)	1.06 ± 0.08	1.16 ± 0.12	1.09 ± 0.07	1.15 ± 0.06	ns						
Fe (µmol/l)	16.18 ± 9.67	22.74 ± 12.22	20.46 ± 4.20	28.08 ± 14.06	ns						



Vol-12 Issue-01 April 2023

				•	JI-14	12 155uc-01 April 2023					
Zn (mmol/l)	14.18 ± 2.20	42.18 ± 7.01	29.62 ± 5.48	29.88 ± 7.35	非非	**	*	÷	ns	ns	ns
Erythrocytes (× 10 ¹² /l)	6.38 ± 0.66	7.07 ± 0.45	6.96 ± 0.30	6.40 ± 1.28	ns						
Hematocrit (%)	26.53 ± 2.72	33.00 ± 1.23	31.94 ± 1.91	29.15 ± 4.78		赤赤	*	ns	ns	ns	ns
Haemoglobin (g/l)	76.25 ± 7.65	92.40 ± 4.77	90.75 ± 5.85	83.63 ± 11.59	**	*	*	ns	ns	ns	ns
Leukocytes (× 10 ⁹ /l)	15.70 ± 3.78	14.94 ± 2.52	14.43 ± 2.10	17.40 ± 6.54	ns						
Lymphocytes (%)	55.50 ± 8.83	65.00 ± 13.40	59.25 ± 6.58	56.50 ± 13.66	ns						
Neutrophils (%)	43.19 ± 8.77	32.60 ± 13.61	39.19 ± 7.40	41.50 ± 14.47	ns						
Monocytes (%)	0.25 ± 0.38	0.60 ± 0.42	0.25 ± 0.46	0.19 ± 0.37	ns						
Eosinophils (%)	0.88 ± 0.74	1.70 ± 0.84	1.31 ± 1.00	1.56 ± 1.27	ns						
Basophils (%)	0.19 ± 0.26	0.10 ± 0.22	0.25 ± 0.27	0.25 ± 0.38	ns						

Control = basal diet containing 110 mg ZnO/kg, ZnO 2.5 = basal diet supplemented with 2.5 g ZnO/kg, ZnO 1.7 + HNa (f) = basal diet supplemented with 1.7 g ZnO and 20.0 g HNa/kg, ZnO 1.7 + HNa (w) = basal diet supplemented with 1.7 g ZnO/kg and drinking water supplemented with 0.2% HNa

 $P \stackrel{<}{<} 0.05$, $P \stackrel{<}{<} 0.01$ described to decrease serum TGC and cholesterol in animals (Samudovska and

Demeterova 2010; Mista et al. 2012; Ozturk et al. 2012). In our study, partial substitution of ZnO by HNa in feed resultedin significantly lower serum TGC in comparison with ZnO 2.5, but this was not observed when HNa was supplemented in water. This variability maybe because of differences in the form and amount of HNa received in feed and water. Neither total cholesterol, nor HDL and LDL were affected by HNa supplementation.

Piglets treated with ZnO and HNa had (ZnO

1.7 + HNa (w)) or tended to have (ZnO 1.7 +HNa(f)) lower urea in serum. A trend towards a de-crease in blood urea nitrogen after HS dietary supplementation was found previously in laying hens (Rath et al. 2006). Serum mineral levels (Ca, P, Mg and Fe) were not affected by ZnO + HNa treatments. HS appear to have metal-chelating ef- fects and were reported to reduce levels of serum minerals in broilers (Rath et al. 2006; Celik et al. 2008; Samudovska and Demeterova 2010) and rab- bits (Mista et al. 2012), although this effect was not observed with HNa in weaned piglets (Trckova et al. 2015). Significantly higher Zn levels in the serum of the ZnO 2.5 group reflected the higher Zn dietary intake in the group.

In the blood of treated piglets, significantly or non-significantly higher HCT and HGB were detected in comparison with Control (Table 2). This

confirms the previous findings that dietary Zn ap- peared to be responsible for an increase of HCT and HGB in pigs (Rupic et al. 1998). The

increase of HCT and HGB was also observed when HA was fed to broilers and rabbits (Ipek et al. 2008; Mista et al. 2012), although no data have as yet been report-ed from pigs. In our previous study, we observed non-significantly higher HCT and HGB when ZnO and HNa were fed to piglets challenged by E. coli (Trckova et al. 2015). Blood cell content was not affected by treatments.

Biomarker of oxidative stress Isoprostanes are specific products of non-enzymatic free radical-catalysed peroxidation of arachidonic acid and bioactive F₂isoprostanes (mainly 8-*iso*-PGF_{2 α}) detectable

in tissues and

body fluids were shown to be the most reliable

in vivo biomarkers of lipid peroxidation and oxida- tive stress in animal and human studies (Kadiiskaet al. 2005; Basu 2007; Basu 2008; Niki 2014). Quantification of F₂-isoprostanes in plasma or se- rum is a specific, reliable and non-invasive method used also for evaluation of various dietary anti- oxidants and other free radical scavengers (Basu 2007; Basu 2008). In our study, significantly lower (P < 0.05) serum

8-iso-PGF_{2a} levels were found in ZnO 1.7 + HNa

(w) (61 pg/ml) and Control (91 pg/ml) in compari-

son with ZnO 2.5 (454 pg/ml) after three weeks



of dietary treatment. The levels in ZnO 1.7 + HNa (f) (132 pg/ml) were non-significantly lower compared to ZnO 2.5.

Zn is an essential trace element important for growth, immunity, and healing. It has also been shown to have antioxidant properties, although the mechanisms are not fully understood. Chronic Zn deprivation in animals generally results in increased free radical production and higher sensitivity tooxidative stress (Powell 2000). Wang et al. (2009) showed that high dietary Zn (3000 mg/kg as ZnO for 15 days) reduced levels of oxidative stress and prevented apoptosis in the jejunum of weaned piglets. Similarly, feeding 1000-2000 mg/kg Zn for 14 days was found to affect the expression of genes involved in reducing oxidative stress in newly weaning pigs (Martinez-Montemayor et al. 2008). Studies in poul- try confirmed that dietary Zn improved antioxidantstatus and augmented the activity of enzymes (sudoi: 10.17221/70/2016-VETMED

peroxide dismutase and glutathione peroxidase) involved in the antioxidant defence system (Sahin et al. 2005; Ma et al. 2011). However, the results from our study do not suggest such an effect. The high pharmacological dose of ZnO fed to weaned piglets for 21 days seemed to negatively influence the oxidative status of piglets

according to the levels of serum 8-*iso*-PGF_{2 α}.

Some recent studies conducted in animals suggest that

prolonged over-supplementation with dietary Zn can promote oxidative stress with adverse effects on growth performance. Our results are in accordance with Pieper et al. (2015) who found that long-term supplementation with high dietary Zn (2425 mg/kgfor 28 days) triggers oxidative stress reactions in the pancreas of weaned piglets. Similarly, Zhao et al. (2014) found that while 20-60 mg/kg of ZnO nanoparticles in broiler diet stimulate antioxidativefunctions, a higher dose (100 mg/kg) can lead to hepatocyte damage and a gradual decrease in total antioxidant capacity in liver tissue. In support of the aforementioned results, Gazaryan et al. (2002) suggested that Zn^{2+} can interfere with mitochon- drial antioxidant production and may

Vol-12 Issue-01 April 2023

stimulate the production of reactive oxygen species.

Our results showed that partial substitution of ZnO by HNa positively affected the oxidative status of piglets as judged by a decline of 8-*iso*-PGF_{2 α}

in serum in both ZnO + HNa-treated groups as

compared with ZnO 2.5. This could be due to the lower dose of ZnO (1.7 mg/kg) which might rather act in an anti-oxidative manner in comparison to high pharmacological doses. Alternatively, the de- crease in the levels of the oxidative stress biomark- er could be the result of the added HNa. Differing effects of HS on oxidative stress and induction of lipid peroxidation in human and animals have been described. HS containing phenolic and polyphenolic compounds can act as an antioxidant, but also as an enzyme (including anti-oxidativeenzymes, e.g. catalase) inhibitor in a time- and concentration-dependent manner (Varanka et al. 1999). Inhibition of these enzymes leads to higher levels of reactive oxygen species (ROS) and development of oxidative stress with subsequent negative effects in the body. However, HS are substances with very complex structures and it cannot be expected that two separate sources of HS will contain identical molecules and exert the same effects on the animal or human body (Vaskova et al. 2011).doi: 10.17221/70/2016-VETMED

Weber et al. (2014) suggested that HS in the diet of young pigs (0.25% for 35 days) may play a role in negating the effects of oxidative stress within the body. The inclusion of HS in the diets of broilers and growing pigs was found to reduce lipid per- oxidation in muscle tissues during storage which had a positive effect on meat quality (Aksu et al. 2005; Bai et al. 2013). The *in vitro* study of Vaskova et al. (2011) showed a decrease in the activity of superoxide dismutase; however, activities of other antioxidant enzymes were not influenced after HS treatment and HS-supported generation of ROS was not observed.

FA profile in diets and blood serum of piglets FA composition in the diets and HNa is



presented in Table 4. HNa contained a low amount of to-tal FA and its supplementation to the basal dietin a dose of 20 g/kg did not alter the FA contentand composition in diet. Differences between the relative content of

On the other hand, Ipek et al. (2008) demon-

strated that high levels of HS dietary tation (480-600 mg/kg) for five weeks increased oxidative stress in Japanese quails, although low levels (360 mg/kg) did not. Also, an in vitro study on endothelial cells showed that HA enhanced the accumulation of intracellular iron resulting in ROS generation and lipid peroxidation when ad- ministered for a long time at high levels (Gau et al. 2001). According to previous studies in humans, longterm exposure to HA and fulvic acids (through drinking water) induces the generation of ROS andpromotes oxidative stress and lipid peroxidationin both a dose- and timedependent manner. It can lead to oxidative DNA and protein damage, increased permeability of cell membranes and pos-sibly cell death, and also to a reduction in the activity of antioxidant enzymes. This mechanism may contribute to the pathogenesis of Blackfoot disease(peripheral vascular disease). Kashin-Beck disease (chronic osteoarthritic disorder), lung

	Lauric	C12	:0 0.29	
Myristic	C14:0	1.37	1.39	5.00
Pentadecanoic	C15:0	0.22	0.23	2.68
Palmitic	C16:0	21.88	21.89	32.51
Cis-7 hexadecenoic 2.93	C16:1 n-9	0.3	30 0.30)
Palmoleic	C16:1	2.11	2.11	1.23
Heptadecanoic	C17:0	0.31	0.31	1.87
Heptadecenoic	C17:1 n-7	0.20	0.20	0.35
Stearic	C18:0	4.46	4.47	11.45
Vaccenic	C18:1 n-11	0.00	0.00	0.76
Oleic	C18:1 n-9	16.50	16.50	13.30
Cis-vaccenic	C18:1 n-7	3.79	3.77	0.01
Linoleic	C18:2 n-6	38.19	38.18	18.39
γ-Linolenic	C18:3 n-6	0.06	0.06	0.07
Linolenic	C18:3 n-3	5.15	5.15	0.64
Nonadecylic	C19:0	0.01	0.01	0.23
Arachidic	C20:0	0.21	0.20	1.30
Eicosenoic	C20:1 n-9	0.47	0.47	0.42
Eicosadienoic	C20:2 n-6	0.69	0.69	0.00

ISSN: 2320-3730

Vol-12 Issue-01 April 2023

individual FA in the diet and serum of control and treated groups are shown in Figure 1. The results reflect the transformation of Table 4. Fatty acid (FA) composition in the diets and sodium humate (HNa)

supplemen-Fatty acids (%) Structure ^{Basal}Basal diet supplemented HNa diet

emphysema and fibrosis and other pathologies (Cheng et al. 1999; Peng et al. 1999; Gau et al. 2001; Ho et al. 2003; Qi et al. 2008). Despite these reports on the effects of longterm supplementation of high levels of HS, in our study, supplemenation with HNa in a doseof 20 g/kg in feed or 0.2% in drinking water for21 days positively influenced the oxidative sta- tus of piglets in comparison with ZnO 2.5 treat-

ment as judged by 8-*iso*-PGF_{2 α} levels in serum.

It should also be taken into account that wide varia-

^{0.29 1.72}



Mead	C20:3 n-9	0.01	0.01	0.06					
Dihomo-y-linolenic	C20:3 n-6	i	0.01	0.01					
0.00									
Arachidonic	C20:4 n-6	5 0.39	0.39	0.00					
Eicosatrienoic	C20:3 n-3	0.03	0.02	0.00					
Eicosapentaenoic	C20:5 n-3	0.73	0.72	0.00					
Behemic	C22:0	0.18	0.18	1.46					
Erucic	C22:1 n-9	0.12	0.12	0.00					
Adrenic	C22:4 n-6	5 0.22	0.22	0.00					
Docosapentaenoic	C22:5 n-3	0.30	0.30	0.00					
Lignoceric	C24:0	0.24	0.24	2.89					
Nervonic	C24:1 n-9	0.06	0.06	0.00					
Docosahexaenoic	C22:6 n-3	0.86	0.86	0.00					
Other FA		0.64	0.65	0.73					
Total FA (nmol/g) 25571.4 25574.7 165.3									

Vol-12 Issue-01 April 2023



FA after digestion of lipids. In the diet, linoleic (LA) was the most abundant FA, followed by palmitic (PA), oleic (OA) and α -linolenic acids (ALA). In blood serum, the content of essential FA (LA and ALA) decreased by about half, while levels of the other n-3 and n-6 long chain polyunsaturated FA (LC-PUFA), mainly eicosapentaenoic (EPA), do- cosapentaenoic (DPA), docosahexaenoic (DHA), dihomo-ylinolenic (DGLA), arachidonic (AA) and ardenic acids, increased in comparison to the diet. It is evident that young growing piglets can efficiently metabolise digested essential FA into physiologically significant n-3 and n-6 LC-PUFA, mainly AA, EPA, DPA and DHA. ZnO 2.5 and both ZnO 1.7 + HNa (f)/(w) treatments significantly modified FA content and com- position in the serum of piglets in comparison withControl (Table 5). Significantly higher total serum FA was found in ZnO 2.5 compare to Control. This finding is in accordance with the significantly higher serum TGC detected in the group. Although studies about the effect of high dietary ZnO on serum FA in weaned piglets are missing, Zhang et al. (2014) reported that Zn participates in cellular lipid me- tabolism as a structural and functional component for some lipid metabolic enzymes and suggested that it plays an important role in FA metabolism.

A pharmacologic dose of ZnO significantly decreased SFA, monounsaturated (MUFA) and poly- unsaturated (PUFA) n-3 and increased PUI 55 n- Diets D Control D ZnO 2.5 D ZnO 1.7 + HNa (f)

sub 50 t are the include pigs, lipid_metabolism, eicc $_{45}$ noids, performance, serum bior $_{40}$ mistry, haematology, and gas chr. $_{35}$ atography/mass spect ometry.



ISSN: 2320-3730

Vol-12 Issue-01 April 2023

in serum compared to Control. In particular, the proportion of myristic, palmoleic, stearic, nonade- cylic, arachidic, vaccenic, OA, cisvaccenic, EPA, DPA, eicosadienoic, DGLA and adrenic acids was decreased, while the content of PA. AA and LA was increased. These results indicate that high ZnO treatment negatively affected the synthesis of somephysiologically significant LC-PUFA, namely n-3 EPA, DPA and n-6 DGLA derived from ALA and LA. The changes in PUFA n-6 and n-3 caused by highZnO treatment resulted in an unfavourable, higher n-6/n-3 ratio compared to the control animals. A different effect on serum FA composition was detected after a lower dose ZnO treatment sup-plemented with HNa. Both ZnO 1.7 + HNa (f)/(w) treatments significantly increased the proportion of SFA and PUFA n-3 and decreased PUFA n-6 in comparison with ZnO 2.5, although the influence of water and feed HNa treatment on individualFA was different. Generally, it can be concluded that the effect of both treatments on individual FA was similar, but the efficiency was different. Partial substitution of the pharmacological dose of ZnO by HNa decreased the PUFA n-6/n-3 ratio in comparison with ZnO 2.5 and the effect was more pronounced (statistically significant), when HNa was applied to drinking water. In contrast to the high ZnO 2.5 treatment, the lower dose of ZnO supplemented by HNa did not have an unfavour- able effect on the metabolism of physiologically ■ ZnO 1.7 + HNa (w)



Vol-12 Issue-01 April 2023





doi: 10.17221/70/2016-VETMED

ISSN: 2320-3730

Vol-12 Issue-01 April 2023

Table 5. Fatty acid (FA) composition in the serum of piglets

	<u> </u>	eatments	<u> </u>	st-hoc_tests								
Fatty acids (%)	Structure		Main									
Control ZnO 2.5 ZnO 1.7 + HNa (w)	ZnO 1.7 +			HNa (f)								
test:2 1:3 1:4 2:	3 2:4 3:4											
Lauric	C12:0	0.041 ± 0.016	0.044 ± 0.014	0.083 ± 0.005	0.088 ± 0.016	非非	ns	**	**	**	**	ns
Myristic	C14:0	0.527 ± 0.038	0.438 ± 0.028	0.617 ± 0.048	0.554 ± 0.061	**	*	**	ns	**	**	ns
Pentadecanoic	C15:0	0.47 ± 0.133	0.429 ± 0.077	0.458 ± 0.123	0.512 ± 0.121	ns						
Palmitic	C16:0	16.64 ± 0.709	19.18 ± 0.309	17.2 ± 0.547	17.41 ± 0.744	非非	**	ns	ns	₩₩.	非非	ns
Cis-7 hexadeceno	ic C16:1 n-9	0.614 ± 0.062	0.556 ± 0.006	0.715 ± 0.016	0.579 ± 0.066	**	ns	ns	ns	**	ns	÷
Palmoleic	C16:1	1.132 ± 0.153	0.874 ± 0.116	1.091 ± 0.133	0.864 ± 0.096	**	**	ns	**	*	ns	*
Heptadecanoic	C17:0	2.0 ± 0.583	1.453 ± 0.299	1.845 ± 0.461	2.642 ± 1.169	*	ns	ns	ns	ns	*	ns
Heptadecenoic	C17:1 n-7	0.405 ± 0.083	0.338 ± 0.067	0.369 ± 0.095	0.455 ± 0.202	ns						
Stearic	C18:0	15.39 ± 1.018	10.63 ± 0.424	16.28 ± 1.719	16.96 ± 1.512	**	**	ns	ns	赤赤	**	ns
Vaccenic	C18:1 n-11	0.132 ± 0.04	0.06 ± 0.015	0.086 ± 0.045	0.105 ± 0.054	*	*	ns	ns	ns	ns	ns
Oleic	C18:1 n-9	11.52 ± 0.344	9.445 ± 0.591	11.74 ± 0.096	10.9 ± 0.808	**	*	ns	ns	\$\$.\$\$	ns	ns
Cis-vaccenic	C18:1 n-7	2.048 ± 0.177	1.459 ± 0.05	2.06 ± 0.046	1.972 ± 0.204	**	*	ns	ns	**	ns	ns
Linoleic	C18:2 n-6	17.86 ± 1.393	24.49 ± 0.938	18.2 ± 1.674	16.8 ± 0.919	非非	赤赤	ns	ns	**	**	ns
γ-Linolenic	C18:3 n-6	0.349 ± 0.034	0.271 ± 0.069	0.359 ± 0.052	0.249 ± 0.048	非非	ns	ns	**	łł	ns	非非
Linolenic	C18:3 n-3	1.354 ± 0.109	1.294 ± 0.07	1.241 ± 0.234	1.271 ± 0.146	ns						
Nonadecylic	C19:0	0.073 ± 0.014	0.043 ± 0.006	0.07 ± 0.015	0.08 ± 0.019	非非	赤赤	ns	ns	łł	**	ns
Arachidic	C20:0	0.3 ± 0.043	0.142 ± 0.015	0.259 ± 0.044	0.208 ± 0.06	非非	赤赤	ns	**	**	ns	ns
Eicosenoic	C20:1 n-9	0.19 ± 0.015	0.157 ± 0.019	0.182 ± 0.027	0.22 ± 0.099	ns						
Eicosadienoic	C20:2 n-6	0.73 ± 0.046	0.452 ± 0.055	0.710 ± 0.074	0.664 ± 0.153	**	**	ns	ns	÷	ns	ns
Mead	C20:3 n-9	0.312 ± 0.088	0.282 ± 0.029	0.346 ± 0.081	0.396 ± 0.126	ns						
Dihomo-γ-linoler	nic C20:3 n-6	1.053 ± 0.109	0.076 ± 0.008	1.15 ± 0.298	1.4 ± 0.312	**	ns	ns	ns	ns	**	ns
Arachidonic	C20:4 n-6	9.706 ± 0.86	13.860 ± 1.148	10.02 ± 0.714	10.17 ± 0.696	**	赤赤	ns	ns	赤赤	**	ns
Eicosatrienoic	C20:3 n-3	0.112 ± 0.043	0.056 ± 0.026	0.051 ± 0.032	0.069 ± 0.038	*	ns	÷	ns	ns	ns	ns
Eicosapentaenoi	c C20:5 n-3	1.544 ± 0.168	0.977 ± 0.113	1.31 ± 0.196	1.26 ± 0.075	**	赤赤	÷	**	赤赤	łł	ns
Behemic	C22:0	0.194 ± 0.022	0.105 ± 0.012	0.174 ± 0.036	0.15 ± 0.04	**	**	ns	ns	<i>4</i> 4	ns	ns
Erucic	C22:1 n-9	0.081 ± 0.027	0.092 ± 0.056	0.048 ± 0.035	0.084 ± 0.053	ns						
Adrenic	C22:4 n-6	3.582 ± 0.576	2.388 ± 0.168	2.838 ± 0.26	3.267 ± 0.637	**	**	ns	ns	ns	*	ns
Docosapentaeno	ic C22:5 n-3	5.429 ± 0.425	3.55 ± 0.304	5.145 ± 1.338	5.088 ± 0.954	**	**	ns	ns	ns	ns	ns
Lignoceric	C24:0	0.339 ± 0.04	0.354 ± 0.026	0.278 ± 0.04	0.21 ± 0.097	**	ns	ns	**	ns	非非	ns
Nervonic	C24:1 n-9	0.719 ± 0.042	0.654 ± 0.05	0.682 ± 0.03	0.474 ± 0.159	**	ns	ns	**	ns	ns	*
Docosahexaenoi	c C22:6 n-3	4.535 ± 0.481	4.698 ± 0.468	4.528 ± 0.395	4.929 ± 0.285	ns						
Other FA		0.059 ± 0.019	0.032 ± 0.019	0.057 ± 0.021	0.066 ± 0.036	ns						
SFA		35.99 ± 1.061	32.83 ± 0.657	37.32 ± 1.538	38.89 ± 0.649	非非	赤赤	ns	非非	杂杂	**	*
MUFA		17.11 ± 0.876	14.75 ± 2.547	16.72 ± 1.058	15.66 ± 1.06	*	÷	ns	ns	ns	ns	ns
PUFA		46.82 ± 0.889	52.39 ± 2.03	45.9 ± 0.971	45.390 ± 1.168	**	**	ns	ns	赤赤	非非	ns
PUFA n-3		13.2 ± 0.594	10.57 ± 0.562	12.28 ± 1.12	12.440 ± 0.878	**	赤赤	ns	ns	*	非非	ns
PUFA n-6		33.31 ± 0.879	41.53 ± 1.502	33.28 ± 1.756	32.550 ± 0.767	**	赤赤	ns	ns	专奏	非非	ns
PUFA n-6/n-3		2.528 ± 0.149	3.931 ± 0.103	2.739 ± 0.397	2.627 ± 0.184	非非	赤赤	ns	ns	ns	*	ns
Total FA (nmol/m	ul)	1118.5 ± 123.6	1435 ± 133.1	1127.4 ± 188.7	1154.8 ± 246.6	*	*	ns	ns	ns	ns	ns
	,							~	~~	~		~

Control = basal diet containing 110 mg ZnO/kg, FA = fatty acids, MUFA = monounsaturated fatty acids, PUFA = polyun -



Vol-12 Issue-01 April 2023

saturated fatty acids, SFA = saturated fatty acids, ZnO 2.5 = basal diet supplemented with 2.5 g ZnO/kg, ZnO 1.7 + HNa (f) = basal diet supplemented with 1.7 g ZnO and 20.0 g HNa/kg, ZnO 1.7 + HNa (w) = basal diet supplemented with 1.7 g ZnO/kg and drinking water supplemented with 0.2% HNa $P \approx 0.05$, $P \approx 0.01$



significant LC-PUFA n-3 and n-6. Levels of most individual PUFA n-3 and n-6 as well as the n-6/n-3ratio in both treatments were similar to Control without treatment. On the other hand, the compo- sition of some individual SFA (lauric, myristic, hep- tadecanoic, arachidic, lignoceric) was significantly higher. The effect on SFA was more pronounced when HNa was applied to drinking water. It is not clear whether the lower dose of ZnO, the presence of HNa, or a combined effect of both was responsible for the obtained results.

To the authors' knowledge, there are currentlyno data available on the effect of HS on serum FA composition in weaned piglets. The results can be partly compared to the findings of Wang etal. (2008), who reported a significant increase in SFA, mainly myristic, palmitic and stearic acids, and USFA in the meat of finishing pigs after 5% HS dietary supplementation. Those authors also detected a significant increase in the USFA : SFA ratio in meat after 10% HS dietary supplementa- tion. Macit et al. (2009) reported an increase in SFA myristic acid and MUFA myristoleic, palmitoleic and heptadecenoic acids in egg volks of layers feda HS-supplemented diet. PUFA n-3 and PUFA n-6 levels in egg yolks remained unchanged by HS di- etary treatments.

In conclusion, the results reported here indicate

that the partial substitution of ZnO by HNa can benefit the performance and health of weaned pig-lets to a similar extent as a high pharmacological dose of ZnO, and at the same time can partially protect against the oxidative stress induced by pro-longed oversupplementation of ZnO in weanedpiglets. Additionally, such a treatment can eliminate the unfavourable effects of high ZnO doses on the n-6/n-3 ratio and the proportion of some physio-logically significant LC-PUFA in serum. Generally, it can be concluded that the effects of both feed andwater HNa supplementation are similar, but that these effects are more pronounced when HNa is applied to drinking water.

ISSN: 2320-3730

Vol-12 Issue-01 April 2023

REFERENCES

Researchers Aksu, Karaoglu, Kaya, Esenbuga, and Macit (2005) examined the effects of dietary humate on the microbiological characteristics, pH, and total bioavailable ribonucleic acid levels in broiler breast and drumstick meats that were vacuum-and aerobically packed. Doi: 10.17221/70/2016-VETMED. Journal of the Science of Food and Agriculture, 85, 1485-1491.

Bai HX, Chang QF, Shi BM, and Shan AS (2013): Fulvic acid's effects on growingfinishing pigs' performance and meat quality. Pages 118–123 in Livestock Science 158. S. Basu (2007): Isoprostanes as a potential new target in the mystery of in vivo oxidative stress assessment. Pages 48–61 of the Scandinavian Journal of Food and Nutrition, volume 51.

The role of F2-isoprostanes in human disease and health: a review of the literature on the subject (Basu, 2008). Redox and Signaling in Antioxidants 10, 1405-1434. A quick technique for extracting and purifying total lipids (Bligh EG, Dyer WJ, 1959). Publication: Canadian Journal of Biochemistry and Physiology, Volume 37, Pages 911–917.

The effects of humic acid and Saccharomyces cerevisiae on performance and metabolic parameters of broiler chickens were studied by Celik K, Uzatici A, and Akin AE (2008). Volume 3, Issue 3, Pages 344–350, Asian Journal of Animal and Veterinary Acknowledgements.

Anemia in Blackfoot disease may be caused by humic acid-mediated oxidative damages to human erythrocytes, according to research published by Cheng ML, Ho HY, Chiu DT, and Lu FJ in 1999. Articles 470–477 in the journal Free Radical Biology and Medicine [27].

EC Regulation 1831/2003 - On September 22,



2003, the European Parliament and Council passed Regulation (EC) No. 1831/2003 on animal nutrition supplements. Citation: EMEA - The European Agency for the Evaluation of Medicinal Products (1999): Humic acids and their sodium salts, Summary report, Official Journal of the European Union, The European Parliament and the Council of the European Union, Brussels, L268/29-L268/43. Committee for Veterinary Me- dicinal Products. The report "Maximum Residue Limits—

Report/2009/11/WC500014416.pdf" may be accessed at this URL:

http://www.ema.europa.eu/docs/en_GB/docu ment_library/.

In 2001, Gau, Yang, Suen, and Lu published: A potential mechanism leading to atherethromhotic veccular dusfunction in

atherothrombotic vascular dysfunction in blackfoot disease is the induction of oxidative stress by humic acid via an increase in intracellular iron. Research Communications in Biochemistry and Physiology 283, 743-749.

Zinc inhibits thiol oxidoreductase activity and promotes reactive oxygen species formation by lipoamide dehydrogenase (Gazaryan IG, Krasnikov BF, Ashby GA, Thorneley RN, Kristal BS, Brown AM, 2002). Biological Chemistry and Biochemistry 277, 10064-10072.

Review of feeding techniques to reduce postweaning diarrhea without utilizing in-feed antimicrobial agents in weaned pigs (Heo JM, Opapeju FO, Pluske JR, Kim JC, Hampson DJ, Nya-choti CM, 2013). Animal Physiology and Nutrition, Volume 97, Issue 2, Pages 207-237.

For their 2003 publication, Ho KJ, Liu TK, Huang TS, and Lu FJ: In addition to promoting lipid peroxidation, humic acid mediates iron release from ferritin. The publication is titled "GETMED" and the DOI is 10.17221/70/2016.

A potential mechanism for humic acid-

ISSN: 2320-3730

Vol-12 Issue-01 April 2023

induced cytotoxicity: oxidation in vitro. Chapters 100-109 of the Archives of Toxicology, volume 77. Humic acid's effects on Japanese quails' laying performance, total antioxidant capacity, and other hematological markers were studied in a 2008 study by Ipek, Avci, Iriadam, Kaplan, and Denek. Journal of Endocrinology 72, 56-60. The use of humic acid compounds in animal husbandry (Islam KMS, Schuhmacher A, Gropp JM, 2005). Section 4, pages 126–134, Pakistan Journal of Nutrition. Clinical Examination of Farm Animals, edited by Jackson and Cockcroft (2002). Volume 320. Oxford: Wiley-Blackwell. The effects of humic compounds on pig growth performance, carcass characteristics, and ammonia emission were studied by Ji F. McGlone JJ, and Kim SW in 2006. Animal Science, Volume 84, Pages 2482–2490. Kadiiska MB, Gladen BC, Baird DD, Germolec D, Graham LB, Parker CE, Nyska A, Wachsman JT, Ames BN, Basu S, Brot N, Fitzgerald GA, Floyd RA, George M, Heinecke JW, Hatch GE, Hensley K, Lawson JA, Marnett LJ, Morrow JD, Murray DM, Plastaras J, Roberts II LJ, Rokach J, Shigenaga MK, Sohal RS, Sun J, Tice RR, Van Thiel DH, Wellner D, Walter PB, Tomer KB, Mason RP, Barrett JC (2005): Biomarkers of oxidative stress study II: are oxidation products of lipids, proteins, and DNA markers of CCl4 poisoning? Vol. 38, Issues 698–710, Free Radical Biology and Medicine.

In a study conducted by Kaevska et al. (2016), the authors examined the impact of sodium humate and zinc oxide on the makeup of the faecal microbiota in weaned pigs as a means of preventing post-weaning diarrhea. Publication: Veterinarni Medicina 61, 328-336.

In 2005, Kang JX and Wang J: Polyunsaturated fatty acid analysis made easier. Journal of Biochemistry 6, 1–4.



Researchers Kim SW, Hulbert LE, Rachuonyo HA, and McGlone JJ (2004) found that weanling pigs consume humic substances that have been mined for their relative iron availability. The article is published in the Asian-Australasian Journal of Animal Sciences and appears on page 1266-1270.

In their 2012 article "Nutrition and Pathology of Weaned Pigs," Kim JC, Hansen CF, Mullan BP, and Pluske JR discuss dietary approaches that promote proper gut barrier function. Research in Animal Nutrition 173, 3-16.

In 2011, Ma, Niu, Feng, Wang, and Feng investigated the effects of zinc glycine chelate on oxidative stress, trace element levels, and intestinal pattern in broilers. Biochemical and Molecular Biology 142, 546–556. Macit M, Celebi S, Esenbuga N, Karaca H

(2009): Layer performance, egg quality, and fatty acid content of egg yolks as a result of dietary humate supplementation. Page 315-319 of the Journal of the Science of Food and Agriculture, volume 89. Gene expression profile in hepatic tissue of newly weaned pigs on pharmacological zinc and phytase enhanced diets was examined in a 2008 study by Martinez-Montemayor MM, Hill GM, Raney NE, Rilington VD, Tempelman RJ, Link JE, Wilkinson CP, Ramos AM, Ernst CW. Molecular Medicine 9, 421. Microflora management in pigs' intestines (Metzler, Bauer, & Mosenthin, 2005). Volume 18, Issue 13, of the Asian-Australasian Journal of Animal Sciences. Pages 1353–1362. An analysis of the effects of a humic-fatty acid preparation on some biochemical and haematological blood parameters in developing rabbits was conducted by Mista et al. (2012). Volume 15, Issue 3, Pages 395–397, Polish Journal of Veterinary Sciences.

Lipid peroxidation biomarkers in clinical samples (Niki E., 2014). Chapters 809–817 of Biochimica et Biophysica Acta 1840. The

ISSN: 2320-3730

Vol-12 Issue-01 April 2023

National Research Council (NRC) published a report on swine nutrient requirements in 1998. National Academies Press, Washington, 10th updated edition, 210 pages. Research conducted by Ozturk et al. (2012) examined the effects of humic substance supplementation on performance, carcass, GI tract, and meat quality features, as well as some blood parameters in broilers. Pages 59-65 of the Journal of the Science of Food and Agriculture, volume 92. The involvement of humic chemicals in drinking water in Kashin-Beck illness in China: a study by Peng A, Wang WH, Wang CX, Wang ZJ, Rui HF, Wang WZ, and Yang ZW (1999). Volume 107, Issue 2, pages 293-296, Journal of Environmental Health. Villodre, Pieper, Martin, and Schunter The authors of the 2015 study are Tudela. Weise, Klopfleisch, Zentek, Einspanier, and Bondzio. Piglet pancreatic zinc accumulation, enzyme activity, and proteomic profiles as a result of a zinc-rich diet. Volume 30, Issue 3, Pages 30-36, Journal of Trace Elements in Medicine and Biology. Zinc oxide for piglets when they were weaned, Poulsen HD (1995). Section A-Animal Science, Acta Agriculturae Scandinavica 45, 159-167. The antioxidant capabilities of zinc (Powell SR, 2000). Volume 130, Issues 1447–1454, Journal of Nutrition. Lung epithelial cell and lining fluid

Lung epithelial cell and lining fluid antioxidant defense damage caused by humic acid, according to Qi, Hartog, and Bast (2008). Journal of Environmental Toxicology and Pharmacology 26, pages 96-101. The effects of humic acid on broiler chickens were studied by Rath NC, Huff WE, and Huff GR in 2006. In: Poultry Science, volume 85, pages 410–414.

Researchers Rupic, Ivandija, Luterotti, Dominis-Kramaric, and Bo-zac (1998) studied the effects of various zinc diets on the plasma proteins and hemostatic parameters of fattening pigs. Publication: Acta Veterinaria



Vol-12 Issue-01 April 2023 Received: April 27, 2016

Accepted after corrections: October 29, 2016

Hungarica 46, pp.114-126. In a 2005 study, Sahin et al. found that heatdistressed quail showed improved performance and antioxidant status when given zinc supplements from either organic or inorganic sources. Proceedings of the Poultry Science Society, 84, 882–887. In 2010, Samudovska and Demeterova investigated the effects of a diet supplemented with salt and natural humic substances.

humate on performance and selected metabolic variables in broiler chickens. Acta Veterinaria Brno 79, 385–393. Trckova M, Lorencova A, Hazova K, Sramkova Zajacova Z (2015): Prophylaxis of post-weaning diarrhoea in piglets by zinc oxide and sodium humate. Veterinarni Medicina

60, 351–360.

Varanka Z, Szegletes T, Szegletes Z, Nemcsok J, AbrahamM (1999): Relationship between the structure of some humic compounds and their inhibitory effects on carp catalase. Bulletin of Environmental Contamination Tox- icology 63, 751–758.

Vaskova J, Velika B, Pilatova M, Kron I, Vasko L (2011): Effects of humic acids in vitro. In Vitro Cellular and Developmental Biology – Animal 47, 376–382.

Wang Q, Chen YJ, Yoo JS, Kim HJ, Cho JH, Kim IH (2008): Effects of supplemental humic substances on growth performance, blood characteristics and meat quality in finishing pigs. Livestock Science 117, 270–274. Wang X, Ou D, Yin J, Wu G, Wang J (2009): Proteomic analysis reveals altered expression of proteins related to glutathione metabolism and apoptosis in the small intestine of zinc oxide-supplemented piglets. Amino Acids 37, 209–218.doi: 10.17221/70/2016-VETMED

Weber TE, van Sambeek DM, Gabler NK, Kerr BJ, Moreland S, Johal S, Edmonds MS (2014): Effects of dietary humic and butyric acid on growth performance and response to lipopolysaccharide in young pigs. Journal of Animal Sci- ence 92, 4172–4179.

Yin J, Wu MM, Xiao H, Ren WK, Duan JL, Yang G, Li TJ, Yin YL (2014): Development of an antioxidant system after early weaning in piglets. Journal of Animal Science 92, 612–619.

Zhang HB, Wang MS, Wang ZS, Zhou AM, Zhang XM, Dong XW, Peng QH (2014): Supplementation dietary zinc levels on growth performance, carcass traits, and intramuscular fat deposition in weaned piglets. Biological Trace Element Research 161, 69–77.

Zhao CY, Tan SX, Xiao XY, Qiu XS, Pan JQ, Tang ZX (2014):Effects of dietary zinc oxide nanoparticles on growth performance and antioxidative status in broilers. Bio- logical Trace Element Research 160, 361–367. Zhu LH, Zhao KL, Chen XL, Xu JX (2012): Impact of weaning and an antioxidant blend on intestinal barrier function and antioxidant status in pigs. Journal of Animal Science 90, 2581–2589.